

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/258129114>

Oxygen Consumption Changes With Yoga Practices A Systematic Review

ARTICLE *in* JOURNAL OF EVIDENCE-BASED COMPLEMENTARY AND ALTERNATIVE MEDICINE · OCTOBER 2013

DOI: 10.1177/2156587213492770

CITATION

1

READS

261

2 AUTHORS:



[Anupama Tyagi](#)

RMIT University

3 PUBLICATIONS 6 CITATIONS

[SEE PROFILE](#)



[Marc Cohen](#)

RMIT University

99 PUBLICATIONS 944 CITATIONS

[SEE PROFILE](#)

Oxygen Consumption Changes with Yoga Practices: A Systematic Review

Anupama Tyagi, MA, PhD(c)¹; Prof Marc Cohen, PhD, MBBS (Hons)¹

RMIT University, Bundoora, Victoria, Australia

Abstract

Oxygen consumption varies with physical and mental activity as well as pathological conditions. Although there is a strong relationship between yoga and metabolic parameters, the relationship between yoga and oxygen consumption has not yet been formally reviewed. This systematic review attempted to include all studies of yoga that also measured oxygen consumption or metabolic rate as an outcome. A total of 58 studies were located involving between 1 and 104 subjects (average 21). The studies were generally of poor methodological quality and demonstrated great heterogeneity with different experimental designs, yoga practices, time periods and small sample sizes. Studies report, yoga practices to have profound metabolic effect producing both increase and decrease in oxygen consumption, ranging from 383% increase with cobra pose to 40% decrease with mediation. Compared to non-practitioners, basal oxygen consumption was reported to be up to 15% less in regular yoga practitioners and regular yoga practice was reported to have a training effect with oxygen consumption during submaximal exercise decreasing by 36% after 3 months. Yoga breathing practices emphasise breathing patterns and retention ratios as well as unilateral-nostril breathing and these factors appear critical in influencing oxygen consumption. A number of studies report extraordinary volitional control over metabolism in advanced yoga practitioners who appear to be able to survive extended periods in airtight pits and exceed the limits of normal human endurance. More rigorous research with standardised practices is required to determine the mechanisms of yoga's metabolic effects and the relevance of yoga practises in different clinical populations.

Keywords: yogic, meditation, pranayama, metabolic rate/cost, energy expenditure

Introduction

Human metabolism is the result of continuous anabolic and catabolic processes that maintain homeostasis and sustain life. Metabolic pathways include a complex network of nutritional, neuronal and humoral inputs that are integrated by the central and autonomic nervous systems through pathways that monitor and maintain physiological functioning. All metabolic processes generate heat and are ultimately dependent on the expenditure of energy via consumption of oxygen, which drives oxidative phosphorylation.

Energy expenditure is directly related to metabolic rate and oxygen consumption and these terms are often used interchangeably. Monitoring oxygen consumption has received a great deal of interest in determining oxygen delivery to tissues, cardiorespiratory function and metabolic response to activity. Assessment of oxygen consumption is used in determining energy requirements for healthy lifestyles, exercise programs, and critically ill patients⁽¹⁻³⁾ and oxygen consumption is reported to increase with adaptation to physiological stress and pathology^(4, 5). The measurement of energy expenditure can be performed via direct calorimetry, which measures heat loss using insulated chambers, or via indirect calorimetry, which directly measures oxygen consumption⁽⁶⁾ through respiratory gas exchange. Direct calorimetry is not frequently used as it is complex, does not accurately measure rapid changes in metabolism and requires significant expertise and elaborate equipment including specially constructed chambers. Indirect calorimetry, is the most commonly technique for measuring energy expenditure and can be used to measure the substrate of metabolism as well as oxygen consumption, which can be expressed in terms of VO_2 (Absolute oxygen consumption), $\text{VO}_2/\text{kg}/\text{min}$ (Relative oxygen consumption), and MET (Metabolic Equivalent Task)^(2, 3, 7).

Oxygen consumption, stress and pathology

oxygen consumption is maximal during intense physical activity and lowest during basal or resting conditions and naturally increases with both psychological and physiological activity, stress and pathology, and higher oxygen consumption appears to correlate to accelerated aging^(4, 5, 8, 9). oxygen consumption has also been found to

increase with activities such as mental arithmetic and playing video games⁽¹⁰⁻¹³⁾ as well as with psychological distress and anxiety⁽¹⁴⁾. A growing body of research further suggests that oxygen consumption is higher in various pathological conditions including, congestive heart failure⁽¹⁵⁾, locomotor impairment⁽¹⁶⁾, HIV⁽¹⁷⁾ and chronic obstructive pulmonary disease⁽¹⁸⁾, and insomnia⁽²⁾, congestive heart failure⁽¹⁹⁾. Oxygen consumption has also been found to increase with features of Metabolic Syndrome including obesity⁽²⁰⁻²²⁾, Type II Diabetes⁽²³⁻²⁶⁾ and hypertension⁽²⁷⁻²⁹⁾.

The measurement of oxygen consumption can provide insights into overall homeostatic balance and response to stress, which are mediated through multiple pathways under the control of the autonomic nervous system and the hypothalamus. The sympathetic nervous system is involved in rapidly mobilising vital physiological functions via sympathetic-adrenal-medullary pathways (SAM) in response to acute stress⁽³⁰⁻³²⁾ which serves to increase oxygen consumption. Repeated or chronic stressful stimuli may lead to changes in the hypothalamic-adrenal-pituitary axis (HPA) leading to a sustained stress response involving cognitive, emotional, endocrine and immune system changes⁽³³⁾. The parasympathetic nervous system provides a counter to the stress response and reduces oxygen consumption by activating the so-called 'relaxation response'⁽³⁴⁾, which serves to reduce physiological arousal and induce a hypometabolic state mediated via enhanced vagal activity⁽³⁵⁾. Such hypometabolic states are suggested to enhance survival in plants and animals by facilitating restorative and repair functions⁽³⁶⁾.

Yoga, stress and metabolism

Mind-body practices that induce relaxation have been traditionally used by people across cultures to improve health and serve as a path for spiritual awakening⁽³⁷⁾. Yoga is an ancient mind-body approach that combines the practice of postures (*asana*), breathing (*pranayama*) and meditation (*dhyana*) with the aim of achieving an effortless state of harmony (*samadhi*).

Yoga postures include both static and dynamic postures that are designed to attune the body to a stable state suitable for meditation. Yoga breathing includes a range of practices such as Bhastrika (bellows breath), *Ujjayi* (victorious breath), *Kapalbhati* (lustrous cranium) and unilateral-nostril breathing, which can be performed at different rates (reported as breath/min) and with different retention periods and patterns that involve either

internal retention (Inspiration:Retention:Expiration (I:R:E)), or external retention (Expiration:Retention:Inspiration (E:R:I)). The yogic state of meditation is characterised by decreased oxygen consumption and cardiovascular activity^(35, 38) and has been shown to elicit the relaxation response⁽³⁴⁾. This meditative state, which is distinct from rest^(39, 40), physical relaxation⁽⁴¹⁾ and sleep⁽⁴²⁾, may be voluntarily induced, even while performing fixed physiological workloads⁽⁴³⁾.

The ability of yoga to induce relaxation and relieve stress has been widely reported⁽⁴⁴⁻⁴⁶⁾ and there are reports of yoga practices reducing acute, chronic and post-traumatic stress. For example yoga is reported to relieve workplace stress⁽⁴⁷⁾, examination stress^(48, 49) and stress-induced inflammation⁽⁵⁰⁾. Yoga practices have also been reported to improve many clinical conditions such as anxiety⁽⁵¹⁻⁵³⁾, depression^(53, 54), negative mood states⁽⁵⁵⁻⁵⁸⁾ and post-traumatic stress disorder (PTSD) symptoms in war veteran,⁽⁵⁹⁻⁶¹⁾ tsunami survivors^(62, 63), hurricane refugees⁽⁶⁴⁾ and flood survivors⁽⁶⁵⁾. Furthermore, two reviews, one involving 35 clinical studies⁽⁶⁶⁾ and the other 8 controlled trials of healthy adults⁽⁶⁷⁾ acknowledge the promising role of yoga in reducing stress. Li et al. 2012 also suggest yoga as a potential adjunct to pharmacologic therapy for patients with stress and anxiety (66). There are further studies to suggest that regular yoga practice reduce physiological and metabolic activity under normal conditions. Compared to non-practitioners, regular yoga practitioners have been found to have lowered resting heart rate⁽⁶⁸⁾, blood pressure⁽⁶⁸⁾ breath rate⁽⁶⁹⁾ and metabolic rate^(70, 71). Yoga has also been found to improve all features of metabolic syndrome including obesity^(72, 73), hyperlipidaemia⁽⁷⁴⁻⁷⁶⁾, hyperglycemia^(75, 77, 78) and hypertension⁽⁷⁹⁻⁸¹⁾, with three separate randomised controlled trials demonstrating benefits of yoga in metabolic syndrome patients⁽⁸²⁻⁸⁴⁾.

While there seems to be a strong relationship between yoga and metabolic parameters, the relationship between yoga and oxygen consumption has not been formally reviewed. The objective of this paper is to systematically review previous research exploring the relationship between yoga and oxygen consumption and explore the impact that different yoga practices have on oxygen consumption in different populations.

Methodology

For this systematic review, a comprehensive search of multiple databases including Scopus, PUBMED, PSYCHINFO, CINAHL, Science Direct database was conducted and a separate search was conducted in Indian medical journals through IndMed which index over 100 prominent Indian scientific journals. Similarly, a search was performed of Yoga Mimamsa, which includes published yoga research literature dating back from 1920 not listed in the above databases. The archives of the International Journal of Yoga were also searched, along with the reference citations from all full text papers identified. The primary search terms included Yoga, yogic, *pranayama*, *yoga nidra*, breathing, relaxation, meditation, Transcendental meditation, *Brahamakumari* meditation, *Raja* Yoga meditation, *Om* meditation, *mantra* meditation, *Sahaj* Yoga meditation, Cyclic meditation and *Kundalini* yoga, *Kriya* yoga and *Sudarshan kriya* along with key words 'oxygen consumption', 'energy expenditure', 'metabolic cost' and 'metabolic rate'.

All studies that had oxygen consumption (either at resting, during yoga intervention or during physical exercise in which yoga included in the intervention) as an outcome were included in the systematic review. The search was performed for articles published up to Dec 2012 and was not otherwise restricted by date or study population. The review included studies that examined a range of yoga practices including asana and/or integrative yoga, breathing, meditation and yogic relaxation practices used either alone or as an integrated practice. The studies were excluded if they were not in English (n=4), unobtainable (n= 5), in press (n=8) or only documented study protocol (n=5). Studies were also excluded if they only involved meditation (religious or non-religious) and relaxation practices that are not directly associated with yoga such as *Zazen/Zen* Buddhist meditation, *Vipassana* Meditation, *Tum-Mo* yoga, *Qigong*, Relaxation Response (RR), Progressive Muscle Relaxation (PMR) and Autogenic Relaxation (AR). However, it was beyond the scope of this systematic review to collect and synthesize clinical outcomes other than oxygen consumption or critically assess the methodological quality of all studies. The selection of relevant studies is shown in Figure 1 and the results, including their statistical significance are noted in the relevant text and tables.

Results

A total of 58 studies of oxygen consumption and yoga practices were extracted (Figure 1). These studies involved between 1 and 104 subjects (average 21) and demonstrated great heterogeneity with many different experimental

designs, yoga practices and time periods. Extracted studies, which were categorized according to the type of intervention (pranayama practice, meditation/relaxation, integrated yoga/asana practice, integrated yoga with physical activity), are presented in Tables 1-4 which also include information about study design.

Of the total studies, 35 studies were published from India (70, 71, 85-117), 15 from USA (118-132), 2 from UK ^(133, 134) and 1 each from, Mexico ⁽¹³⁵⁾, New Zealand ⁽¹³⁶⁾, Thailand ⁽¹³⁷⁾, Brazil ⁽¹³⁸⁾, Japan ⁽¹³⁹⁾ and Sweden ⁽¹⁴⁰⁾. Most studies reported assessing direct measurement of respired gases for measuring oxygen consumption using indirect calorimetry techniques, whether through open circuit, closed circuit, bag system or respiratory chamber method. Some studies derived the oxygen consumption through standard equations such as oxygen consumption was predicted through regression equation with the measures of heart rate and oxygen consumption of submaximal exercise ⁽⁹⁴⁾, VO₂ max was predicted through achieved workload and using standard formula from American college of sports and medicine ^(116, 130). Oxygen consumption was reported to both increase and decrease with different yoga practices. Increases in oxygen consumption ranged from 7.7% with *Ujjayi* breathing to 383% during cobra pose (Table 1 & 3). Studies also report decreases in oxygen consumption with slow yoga breathing techniques and meditation practices ranging from a 3.7% decrease during Om meditation to a 40% decrease in an advanced yogi during meditation in an air-tight pit (Table 2). Basal oxygen consumption is also reported to be up to 15% less in regular yoga practitioners compared to non-practitioners and oxygen consumption during submaximal exercise is reported to decrease by 36% after 3 months of regular yoga practice (Table 4).

Pranayama Practices and Oxygen Consumption

Table 1 summarises 16 *pranayama* (yogic breathing) studies that include a total of 143 participants and report wide variations in oxygen consumption. While oxygen consumption was seen to increase with most breathing practices performed at both fast (232 breath/min) and slow (1 breath/min) rates (Table 1), a decrease in oxygen consumption from rest was also seen in some slow breathing practices. The highest increase in oxygen consumption was seen with extremely rapid *Bhastrika* breathing, which involves rapid, forced thoracic inhalation and exhalation. When *Bhastrika* was performed at a rate of 232 breath/min by 3 advanced practitioners oxygen consumption was reported to increase by 208% ⁽¹⁴⁰⁾ and increases in oxygen consumption of 30%, 24%, 22%, 17%

and 15% are reported with *Bhastrika* performed at different rates and retention periods ^(88, 90, 118, 119). Increases in oxygen consumption of 12% ⁽¹¹⁹⁾ to 50% ⁽⁸⁷⁾ are also reported with *Kapalbhati* breathing, which involves forced rapid exhalation. Unilateral nostril breathings (alternate nostril breathing, right nostril breathing and left nostril breathing) are reported to increase oxygen consumption with a 150% increase during alternate nostril breathing ⁽⁹⁴⁾ and increases of 37% ⁽⁹⁶⁾ to 18% ^(93, 96) reported immediately after alternate nostril breathing (ANB), right nostril breathing (RNB) and left nostril breathing (LNB) practices.

Oxygen consumption is also reported to increase with some slow yoga breathing. *Ujjayi* breathing, which involves controlled slow, deep breathing with long inhalation and exhalation and gentle contraction of the glottis creating a soft snoring sound ⁽¹⁴¹⁾, has been consistently reported to increase oxygen consumption, even at extremely slow rates. An increase of 10% is reported in a single advanced practitioner while practicing *Ujjayi* at a rate of 1 breath/min ⁽¹³⁹⁾, while further studies report increases in oxygen consumption of 25% and 52% during *Ujjayi* with a 40 second retention (rate of 1.26 breath/min) ⁽¹¹⁹⁾ or with I:R:E ratio of 1:1:1 ⁽⁹²⁾. An increase in oxygen consumption was also reported with *Ujjayi* performed at different altitudes with a 16% greater oxygen consumption observed in a single practitioner at 3200m elevation practicing *Ujjayi* breathing at 3 breath/min compared to practicing *Ujjayi* breathing at 520m elevation at 1.5 breath/min ⁽⁸⁶⁾. An increase in oxygen consumption to 17% has also been reported in advance yoga practitioners during slow paced breathing with I:R:E ratio of 1:4:2 ⁽¹³⁸⁾.

Only 4 studies (Table 1) report decreases in oxygen consumption with pranayama. A decrease in oxygen consumption of 4%, 21% and 19% is reported during slow *Ujjayi* breathing at rates of 2 breath/min ⁽⁹⁰⁾, 1.4 breath/min ⁽⁹¹⁾ or with a I:R:E ratio of 1:4:4 ⁽⁹²⁾. A decrease in oxygen consumption of 16% is also reported during *Bhastrika* breathing at 12 breath/min ⁽⁹⁵⁾.

Yoga Meditation, Relaxation Practices and Oxygen Consumption

Table 2 summaries 15 studies with a total of 310 participants that consistently report reduced oxygen consumption during different meditation and relaxation practices. Two studies of yogic relaxation practices report 25.2% and 23% reductions in oxygen consumption compared to rest ^(100, 101). Transcendental meditation is also reported to

produce reductions of oxygen consumption from rest with 3 separate studies reporting reductions of 20%, 17% and 5% ⁽¹²⁰⁻¹²²⁾. Reductions in oxygen consumption from rest of 15% and 3.7% are further reported during 2-3 minutes of meditation ⁽⁹⁵⁾.

Studies comparing meditation with non-yogic relaxation techniques report modest or no difference between interventions. Four studies report no difference in oxygen consumption between groups practicing Transcendental and those practicing a control relaxation intervention ^(123, 124, 134, 136), while a further study reports no significant reduction in oxygen consumption from baseline rest during either after *Om* meditation or relaxed sitting, despite reported reductions in heart rate and increases in galvanic skin response⁽¹¹⁷⁾.

Among the studies reporting reductions in oxygen consumption, the most dramatic reductions were seen in two studies involving advance yoga practitioners, with one study reporting reductions in oxygen consumption of 40% below rest during a 4 hour stay in an air tight subterranean chamber ⁽⁹⁹⁾ and another study reporting reductions of 32% and 37% below rest during two separate 10 hour stays in an air tight box ⁽⁹⁷⁾. Reductions in oxygen consumption of around 35% below rest are also reported during meditation in a group of experienced yogis (n=9), ⁽¹³⁸⁾. An early study with 3 advanced yoga practitioners further reports that during a prolonged stay in an air tight pit, advanced meditators could tolerate ambient O₂ levels of 12.2% and CO₂ levels of 7.3% ⁽⁹⁸⁾.

Asana/Integrated Yoga Practices and Oxygen consumption

Table 3 presents 13 studies with a total of 272 subjects that consistently report increases in oxygen consumption with different yoga *asanas* (postures). The most dramatic increase was seen in a group of 21 male practitioners who experienced a 383% increase in oxygen consumption while performing cobra pose ⁽¹⁰⁴⁾. Increases in oxygen consumption were also reported with warrior III pose (300%) ⁽¹²⁵⁾, plough pose 2 (160%) ⁽⁹⁵⁾, Hero pose (159%) ⁽¹⁰³⁾, headstand pose (68%) (85) and accomplished pose (27%) ⁽¹⁰²⁾.

Over the course of a yoga session oxygen consumption has been reported to increase by 100% with *Ashtanga* yoga ⁽¹²⁶⁾, 114% with *Hatha* yoga ⁽¹³¹⁾, 133% with Thai yoga ⁽¹³⁷⁾ and 144% with *Iyengar* yoga⁽¹²⁵⁾. Three studies have examined oxygen consumption during Sun Salutation (a dynamic sequence of 12 postures) and report that oxygen

consumption increased 205% above resting levels ⁽¹⁰⁴⁾ and 25% ⁽¹²⁶⁾ and 81% ⁽¹³¹⁾ above the levels during static postures.

The reported increases in oxygen consumption seen with yoga practices are less than observed with maximal or submaximal exercise. oxygen consumption during Thai yoga is reported to be 35.5% of VO_{2max} ⁽¹³⁷⁾ and *Vinyasa* yoga, 50% ⁽¹²⁷⁾, bow posture 26.5% and *Shavasana* (supine pose), 9.9% ⁽⁹⁵⁾ of VO_{2max} . Similarly *Iyengar*, *Ashtanga* and *Hatha* yoga sequences have been shown to be of lower intensity than sub-maximal exercise, having oxygen consumption that is 26%, 33% and 54% lower than oxygen consumption during treadmill walking at 4mph ⁽¹³²⁾, 3 mph ⁽¹²⁶⁾ or 3.5mph ⁽¹³¹⁾ respectively.

While oxygen consumption is reported to increase during a yoga session, there are reports that oxygen consumption may fall below pre-session levels immediately after certain practices. During Cyclic meditation, which involves a series of postural sequences interspersed with periods of relaxation, oxygen consumption is reported to increase by up to 55% during the active phase and then fall to 19% below pre-session levels in the immediate post session period ⁽¹⁰⁶⁾. Similar results are reported in a further study which reports a 32% decrease in oxygen consumption immediately after Cyclic mediation ⁽¹⁰⁵⁾.

Regular Yoga Practice, Physical Activity and Oxygen Consumption

Table 4 presents 16 studies involving 516 participants that measured oxygen consumption at rest or during physical activity (sub maximal and maximal) after 1 month to 24 month of integrated yoga practice (including *asana*, *pranayama* and relaxation) along with two studies comparing oxygen consumption at rest in yoga and non-yoga practitioners ^(70, 71) and one study comparing oxygen consumption between groups who regularly practiced lotus posture and groups of regular exercisers or healthy sedentary subjects ⁽¹⁰⁷⁾.

Most of these studies report regular yoga practice leads to progressive reductions in oxygen consumption over time. In a 3 months cohort study, yoga practice was found to reduce oxygen consumption during submaximal exercise by 36% compared to baseline levels ⁽¹¹⁰⁾. A randomized trial involving male soldiers found that 6 months yoga practice (n=15) reduced oxygen consumption during submaximal exercise by 5.7% (P<0.05) compared to no

change in a physical training group (n=15)⁽¹¹⁴⁾, while a non-randomised study reports that 12 months of regular yoga practice with regular sports activity improved submaximal work efficiency in athletes with 51% greater work output per litre of oxygen consumed, compared to no change in regular sports activity group⁽¹¹²⁾.

VO_{2max} was also reported to increase with regular yoga practice ranging from 6 weeks to 6 months in diverse populations. A 3% increase in VO_{2max} is reported in the cohort of middle aged yoga practitioners who practiced intensive yoga for 11 weeks⁽¹⁴²⁾ and 7% increase in VO_{2max} in cohort of yoga navies who practiced integrated yoga for 8 weeks⁽¹²⁸⁾. Similarly up to 7% increment of VO_{2max} is reported in randomized trial of 6 months in male soldiers with integrated yoga (n= 17) compared to no change in a physical training group (n=11)⁽¹¹⁵⁾ and a 13% (P<01) increase in VO_{2max} is reported in elderly subjects in randomised trial after 6 weeks of yoga with practice (n=20), similar to significant increase with aerobic training (n=20)⁽¹³³⁾.

Increases in VO_{2max} of around 17% are also reported after yoga practice in two cohort studies including a 6 week study of healthy subjects (n=17)⁽¹¹⁶⁾, and an 11 week study of elderly yoga practitioners (n= 9)⁽¹⁴²⁾. Similar increases in oxygen consumption are reported in an 8 week randomised controlled trial of patients with congestive heart failure who practiced yoga (n=9), compared to no change in a standard medical therapy group (n=10)⁽¹³⁰⁾. A further cohort study of female physical trainers found that one month of yoga practice led to 14% greater maximal work efficiency⁽¹¹¹⁾. Maximal work efficiency was also seen to improve in non-randomised controlled trial by 34% in athletes after 24 months of regular yoga practice compared to a control group practicing physical exercise⁽¹¹²⁾.

Not all the studies report improvement in oxygen consumption or work efficiency with regular yoga practice. A 12 month randomised study reports no change in oxygen consumption during submaximal exercise in either a yoga or aerobic training group⁽¹¹³⁾. In another randomised study no change in VO_{2max} is reported after 8 weeks yoga practice group (n=10) compared to no-intervention control group (n=11)⁽¹²⁹⁾. Similarly, two 3 months cohort studies report no change in oxygen consumption at rest after regular yoga practice^(109, 110) and similar results are reported in a 12 month randomised controlled trial⁽¹¹³⁾. In contrast to most of the above mentioned studies, one small cohort study reported increased oxygen consumption during submaximal exercise after 6 months of regular yoga practice in healthy subjects despite an observed reduction in resting core body temperature⁽¹⁰⁸⁾.

When examining oxygen consumption at rest, two studies report basal oxygen consumption to be significantly less in regular yoga practitioners compared to non-yoga practitioners. One study⁽⁷⁰⁾ reports that regular yoga practitioners had basal metabolic rate (BMR) 13% less than predicted based on the FAO/WHO/UNU equation⁽¹⁴³⁾ and that oxygen consumption during basal conditions was significantly less in regular yoga practitioners compared to non-yoga practitioners. Similar results were reported in a second study, which report that regular yoga practitioners had basal metabolic rate that was 17.8% less than non-yoga practitioners⁽⁷¹⁾.

Discussion

Studies published to date suggest that yoga practices can have profound metabolic effects producing both significant increases and decreases in oxygen consumption. Like other physical activity, physical yoga postures can increase oxygen consumption dramatically, yet yoga practices do not involve maximal exertion. For example, dynamic postures such as cobra pose are reported to increase oxygen consumption by 383% or around 1220ml/min, which is less than half that produced with maximal exercise in the average untrained healthy male⁽³⁾. The most dramatic change seen with yoga is reduction of oxygen consumption with reports of yoga practices down regulating the sympathetic nervous system and producing modest reductions in oxygen consumption comparable to practices such as progressive muscle relaxation, closed eyes relaxation and listening to music^(123, 124, 134, 136) as well as reports of reductions of dramatic reductions up to 40%⁽⁹⁹⁾. This suggests that yoga may down-regulate the hypothalamic-pituitary-adrenal (HPA) axis and the sympathetic activity and therefore promote relaxation and stress relief.

Regular yoga practice also appears to have a training effect, with regular yoga practitioners consistently showing significant reductions in oxygen consumption during normal physical activity compared to non-yoga practitioners. Thus, unlike other physical training, which generally increases resting metabolic rate^(144, 145), regular yoga practice is reported to decrease resting oxygen consumption to levels lower than predicted by the FAO /WHO/UNU equation⁽⁷⁰⁾. This may be due to regular physical training producing an increase of muscle mass which requires greater oxygen consumption supply at rest, whereas yoga training may instead increase efficiency of mitochondrial oxidative phosphorylation and reduce O₂ demand.

Yoga practises are also reported to shift lactate threshold (anaerobic threshold) and improve work efficiency indicating aerobic capacity and reduced muscle fatigue to a greater degree compared to physical activity ⁽¹¹²⁾ and these results are supported by randomised crossover trial documenting reduction in blood lactate, heart rate and BP with regular yoga practice ⁽¹⁴⁶⁾.

A recent review of yoga and exercise found that yoga may be as effective as, or better than exercise at improving a variety of health-related outcome measures in both healthy and diseased populations. ⁽¹⁴⁷⁾ Despite multiple studies demonstrating the benefits of yoga in various clinical conditions, only one small study examined the effects of yoga and oxygen consumption in a clinical population. This study reported increased aerobic capacity (VO₂ max) in patients with congestive heart failure after practicing yoga postures, breathing techniques and meditation over a period of 8 weeks ⁽¹³⁰⁾. Previous research also suggests that instruction on respiration and relaxation in addition to physical exercise enhances respiratory sinus arrhythmia and slows heart rate and breath rate in myocardial infarction patients during rehabilitation ⁽¹⁴⁸⁾ and that slow rhythmic respiration can be used as a therapeutic tool for anxiety ⁽¹⁴⁹⁾, hypertension ^(150, 151), and asthma ⁽¹⁵²⁾. Due to the wide variety of yoga practices and styles, further research is required to determine the most appropriate practices for different clinical conditions. Typical yoga sessions of different styles appear to differ in exercise stimulus resulting in varied increase in oxygen consumption ^(125, 126, 131, 137) with profound increase reported during dynamic posture sequences compared to static posture sequences ^(126, 131). Different yoga practices and styles however, are likely to have different health and fitness benefits ^(153, 154).

It appears that breath rate and retention periods are critical in determining oxygen consumption and that yoga practitioners are able to vary their breath rate widely with reported breath rates ranging from 1 breath/min to over 230 breath/min. Oxygen consumption is also reported to paradoxically increase by up to 10% despite breath rates of only 1 breath/min. The most profound changes in oxygen consumption with breathing techniques are seen in advanced yoga practitioners who are reported to increase their oxygen consumption by 208% and their CO₂ exhalation by 395% when performing *Bhastrika* breathing at 232 breath/min, or decrease their oxygen consumption by 16% when performing the same type of breathing at 12 breath/min. Similarly, altering the

retention period during *Ujjayi* breathing is reported to either increase oxygen consumption by up to 52% when performed with a short retention period with I:R:E of 1:1:1 or decrease by 19% when the same type of breathing is performed with a longer retention period of I:R:E of 1:4:4. Ultradian rhythms in nasal cycles and unilateral-nostril breathing practices may also influence oxygen consumption with alternate nostril breathing being reported to increase oxygen consumption by up to 150%.⁽⁹⁴⁾ Advanced yoga practitioners appear to be able to exert extraordinary conscious manipulation of their metabolic and autonomic functions^(155, 156), with reports of yogis being able to tolerate ambient CO₂ levels of more than 7% and O₂ levels less than 12%⁽⁹⁸⁾. There are further reports of advanced yogis being able to reduce oxygen consumption by 40% while meditating in an airtight pit⁽⁹⁹⁾ and survive 8 days in an airtight pit with an unrecordable ECG⁽¹⁵⁷⁾. These reports appear inexplicable, yet are similar to reports of advanced Zen meditators being able to decrease oxygen consumption up to 20% along with dramatic decrease in respiratory rate to 1.5 to 2 breath/min during Zazen meditation, Tum-mo meditators being able to increase or decrease their oxygen consumption by over 60% during seated meditation⁽¹⁵⁸⁾, or reports of modern free divers being able to hold their breath for over 10 minutes while diving to depths of over 200m⁽¹⁵⁹⁾. So far, these extreme feats of metabolic control are poorly documented and limited to single case studies or small cohorts. They therefore require further investigation and documentation as they may provide clues about extending the limits of human endurance and metabolic control.

This review suggests that yoga can have profound metabolic effects with a consistent picture emerging from experimental, cohort, non-randomized and randomized controlled trial studies. Yet most of the studies are of poor methodological quality and do not provide adequate reporting of the study design, study population, yoga practices, methods of measurements or statistical methods. Furthermore, most studies were performed in India (n=35) and included only small numbers of adult male yoga practitioners without matched comparison groups. Furthermore, there are 2 randomized controlled trials of healthy people that report no change in oxygen consumption with yoga despite significant changes in other physiological measures. Of these a controlled trial (n=10) reported significant improvements in flexibility with yoga but no change in maximal aerobic capacity⁽¹²⁹⁾, while another controlled trial (n=18) reported improvements in respiratory variables and breath hold time but no change in oxygen consumption during submaximal exercise with yoga⁽¹¹³⁾. A further cohort study (n=10) reported

significant improvements in biochemical and anthropometric parameters after 3 months of yoga practice but did not find any change in oxygen consumption⁽¹⁰⁹⁾.

The small sample sizes, variable practices, and limited, non-clinical populations involved in the reviewed studies make it difficult to generalise results to wider populations or make definitive statements about specific practices. Thus more rigorous studies with larger samples and standardised practices are required to determine the role of yoga in modulating oxygen consumption and determine if the reported results can be reproduced in non-Indian, female, adolescent and non yoga-practicing populations as well as in different clinical conditions. The reports of advanced yogis performing extraordinary feats also warrant further investigation using modern equipment and research methodologies.

Conclusion

Research to date on yoga and metabolism includes many heterogeneous yoga practices in studies of poor methodological quality. This research suggests that yoga practices can produce dramatic changes in oxygen consumption and metabolism and that regular yoga practice may lead to reduced resting metabolic rate. Research further suggests that different yoga postures and breathing practices, which involve the control of respiratory rate and retention periods, may produce markedly different metabolic effects with reductions in oxygen consumption being more dramatic than increases. The extraordinary volitional control over autonomic functions and remarkable feats of metabolic endurance demonstrated by advanced yoga practitioners warrant further investigation and further more rigorous research on standardised practise is required to determine the relevance of yoga practices in various clinical conditions.

Disclosures

Acknowledgement

The work for this article was all performed at RMIT University. The authors would like to acknowledge the assistance of RMIT University biomedical librarian Savita Hazari for her help in conducting the searches and sourcing and obtaining articles.

Author Contribution

Anupama Tyagi was responsible for conducting the literature searches, preparing the tables and writing the first draft of the article. Marc Cohen was responsible for conceiving the article, categorising the papers and assisting in writing the article and reviewing drafts.

Declaration of Conflicting Interests

There are no conflicting interests.

Funding

This article was prepared as part of Anupama Tyagi's PhD research. No external funds or grants were sought or provided.

Ethical Approval

As this article represents a systematic review of literature and no human or animal experimentation, no ethics review was sought or required.




Table 1: Summary Of Studies Reporting Changes in Oxygen Consumption (OC) With Pranayama Practice(s)

Study Reference	Population	Study Design	Intervention	Comparators	Metabolic Measures	Cardio-respiratory and Other Measures
Miyamura, Nishimura et al. 2002 ⁽¹³⁹⁾	Advanced male yoga practitioner (n=1);	Single practice on a single occasion	<i>Ujjayi</i> breathing at 1 breath/min	<i>Pranayama</i> versus <i>post-pranayama</i>	↑ of 10% in OC during <i>Ujjayi</i>	HR ⁺ during breathing session 9% higher compared to post.
Miles and Behanan 1934 ⁽¹¹⁸⁾	Male yoga practitioner (n=1);	Multiple practices on a single occasion	<i>Ujjayi</i> , <i>Kapalbhati</i> and <i>Bhastrika</i> breathing	Sitting and reclined postures versus <i>Pranayama</i>	↑ of 33%, 35% and 30% in OC of during <i>pranayama</i> compared to sitting and 39%, 41% & 36% compared to reclined	
Miles 1964 ⁽¹¹⁹⁾	Male yoga practitioner (n=1);	Multiple practices on multiple occasions	<i>Ujjayi</i> (40 seconds retention at 1.26 breath/min); <i>Kapalbhati</i> (12.5 & 80 breath/min) <i>Bhastrika</i> (21 & 1.3 breath/min);	Baseline versus <i>pranayama</i> practices	↑ of 25% in OC during <i>Ujjayi</i> , 12% during <i>Kapabhati</i> and 19% during <i>Bhastrika</i>	
Rao 1968 ⁽⁸⁶⁾	Male yoga practitioner (n=1);	Single practice on two occasions	<i>Ujjayi</i> breathing at two different altitude of 520 meter and 3800 meter	Baseline versus <i>Ujjayi</i> breathing at low altitude; Baseline breathing versus <i>Ujjayi</i> at high altitude	↑ of 7.7% in OC during <i>Ujjayi</i> with breath rate 1.5 breath/min at low altitude; ↑ of 9.9% in OC during <i>Ujjayi</i> at breath rate 3 breath/min at high altitude; OC during <i>Ujjayi</i> at high altitude was 16% higher compared to lower altitude	

Karambelkar 1982 (87)	Male yoga practitioners (n=8);	Multiple practices on a single occasion	<i>Kapalbhati</i> at 120 breath/min and Hyperventilation breathing at 26 breath/min	Baseline breathing versus <i>Kapalbhati</i> breathing and hyper-ventilative breathing	↑ of 50% in OC and ↑ of 33% in CO ₂ [†] exhalation during <i>Kapalbhati</i> ; ↑ of 133% in OC and 379% in CO ₂ exhalation during hyper ventilative breathing;	↑ of 209% in MV [§] & ↓ of 63% in VT during <i>Kapalbhati</i> ; ↑ of 538% in MV and ↑ of 250% in VT during hyper-ventilative breathing
Karambelkar 1982 (88)	Male yoga practitioners (n=3);	Multiple practices on a single occasion	<i>Bhastrika</i> with internal retention (I:R:E [¶] - 8:32:16) and external retention (E:R:I ^{**} - 3:20:10);	Baseline versus <i>Bhastrika</i> breathing with internal retention and external retention	Internal Retention: ↑ of 15% in OC and ↑ of 13% in CO ₂ exhalation during <i>Bhastrika</i> ; External retention: ↑ of 17% in OC and ↑ of 32% in CO ₂ exhalation during <i>Bastrika</i> ;	
Frostell, Pande et al. 1983 (140)	Experienced male yoga practitioners (n=3);	Single practice on a single occasion	<i>Bhastrika</i> at 232 breath/min	<i>Baseline</i> versus <i>pranayama</i>	↑ of 208% in OC and ↑ of 395% in CO ₂ exhalation during <i>Bhastrika</i> ;	↑ of 30 BPM ^{††} (47%) in HR ↑ of 88.4 l/min (15 fold increase) in MV ; ↑ of 65% in CO ^{††}
Karambelkar 1988 (89)	Male yoga practitioners (n=7);	Single practice on a single occasion	<i>Kapalbhati</i> at 120 breath/min	<i>Pranayama</i> practices versus baseline	↑ of 51% in OC and ↑ of 34% in CO ₂ exhalation during <i>Kapalbhati</i> ;	↑ of 219% in MV during breathing;
Karambelkar 1983 (90)	Male yoga practitioners (n=3);	Multiple practices on a single occasion	<i>Bhastrika</i> with external retention (E:R:I- 3:20:10) and <i>Ujjayi</i> with external retention	<i>Pranayama</i> practices versus baseline	↑ of 24% in OC and ↑ of 32% in CO ₂ exhalation during <i>Bhastrika</i> ; ↓ of 4% in OC and ↓ of 7% in CO ₂	

Karambelkar 1983 (91)	Regular and beginner male yoga practitioners (n=9);	Single practice on a single occasion	(E:R:I- 6:12:12); <i>Ujjayi</i> breathing with internal (I:R:E- 8:32:16)	<i>Pranayama</i> versus baseline	exhalation during <i>Ujjayi</i> ; Significant ↓ of 21% in OC and ↑ of 34% in OC was observed during the same practices, ↓OC was only seen in the regular yoga practitioners.	
Danucalov, Simões et al. 2008 (138)	Experienced yoga practitioners with >3 years' experience (n=9);	Multiple practices on a single occasion	Slow paced <i>pranayama</i> with extended period of retention (Internal retention- I:R:E -1:4:2) and meditation; Each phase of 30 minutes;	<i>Pranayama</i> versus meditation and baseline	↑ of 20% in OC and ↑ of 25% in CO ₂ exhalation during <i>pranayama</i> compared to baseline; ↑ of 84.6% in OC during <i>pranayama</i> compared to meditation	↓ of 35% in OC during meditation compared to baseline.
Telles and Desiraju 1991 (92)	Male yoga practitioners Short breath retention group (n=5); long breath retention group (n=10);	Multiple practices on a single occasion	<i>Ujjayi</i> breathing with 2 different internal retention periods: Short retention (I:R:E- 1:4:4), Long retention (I:R:E- 1:1:1)	Baseline versus <i>Pranayama</i>	↑ of 52% in OC (P<05) during <i>pranayama</i> with short retention (1:1:1); ↓ of 19% in OC (P<025) during <i>pranayama</i> with long retention (1:4:4);	
Telles, Nagarathna et al. 1996 (93)	Male yoga practitioners (n=12);	Multiple practices on two occasions	RNB ^{§§} session and normal breathing session; (Each session of 45 minutes on different days)	Baseline breathing versus Post RNB session and post NB session	↑ of 18% in OC (P<05) after RNB; No significant change after Normal breathing compared to baseline;	↑ of 9.3% SBP (P<05) after RNB; ↓ of 60% in GSR ^{¶¶} after RNB;
Prasad, Venkata	Male Yoga	Multiple	ANB ^{***} for 30 minutes, Treadmill	Resting, field walk and	↑ of 150% in OC (P<01) during	

Ramana et al. 2001 (94)	practitioners with >3 years' experience (n=12);	practices on a single occasion	walk at 3km/hr (1.9 mph) for 30 minutes and field walk 1.5 km/30 min;	treadmill walk versus ANB	ANB compared to resting state; OC during ANB 19.6% (P<05) lower compared to field walk and 37.5%(P<01) lower compared to treadmill walk;
Ray, Pathak et al. 2011 (95)	Male yoga practitioners with >6 years' experience (n=20);	Multiple practices on two occasions	<i>Hatha</i> Yoga session - comprising variety of yoga static postures interspersed with <i>Shavasana</i> , <i>pranayamas</i> and meditation practices; VO _{2max} session; (Each session on different days)	Sitting rest (<i>Sukhasana</i>) versus each individual <i>pranayama</i> versus rest sitting	↓ of 16% in OC during <i>Bhastrika</i> ; ↑ of 65% , 61%,33% (Ps<05) in OC during Raven Beak (<i>Kaki Mudra</i>) breathing (P<05), 61% during I breathing (P<05), 103ml/min (33%) during <i>Kapalbhati</i> (P<05);
Telles, Nagarathna et al. 1994 (96)	Male yoga practitioners (n=48);	4-weeks regular practice in multiple groups	Random assignment to either RNB (n=12), LNB ⁺⁺⁺ (n=12) or ANB (n=24); Each assigned breathing 4 times daily for 4 weeks;	Post <i>pranayama</i> intervention for versus pre intervention	↑ of 37% in OC (P<05) post RNB, ↓ in body weight after 1 month of <i>pranayama</i> ; ↑ of 9.6% & 7% in HR (Ps<001) post RNB and ANB respectively; ↑ of 150% in GSR (P<05) post LNB ;

Table 1: Summary of studies reporting changes in OC with pranayama practice(s)

Abbreviations used: OC*- oxygen consumption; HR[†] – heart rate; CO₂[‡]- carbon dioxide; MV[§] –minute ventilation; VT^{||} –tidal volume; I:R:E[¶] – Inspiration:Retention:Expiration;

E:R:I^{**} - Expiration:Retention:Inspiration; BPM⁺⁺ - beats per minute; CO^{‡‡} - cardiac output; RNB^{§§} – right nostril breathing; SBP^{|||} - systolic blood pressure; GSR^{¶¶} - galvanic skin resistance; ANB^{***} - alternate nostril breathing; LNB⁺⁺⁺ – left nostril breathing

Table 2: Summary Of Studies Reporting Changes in Studies Yoga meditation Oxygen Consumption (OC*) With Meditaion/ Relaxation Practice(s)

Study Reference	Population	Study Design	Intervention	Comparators	Metabolic Measures	Cardio-respiratory and Other Measures
Anand, Chhina et al. 1961 (97)	Experienced male yoga practitioner (n=1);	Single practice on two occasions	Stay in air tight box during two different days of 10 hours each	Baseline (basal) versus stay in box	↓ of 37.4% and 32% in OC during two different sessions.	↓ in HR [†] up to 25BPM [‡] during session; HR rose only when ambient O ₂ declined to 15% and CO ₂ [§] reached to 5% in the pit; OC declined to 50% of BMR (19.5l/hr) on one occasion:
Karambelkar, Vinekar et al. 1968 (98)	Experienced male yoga practitioners (n=4);	Single practice on a single occasion	Stay in air tight pit >12 hours and up to 18 hours	Baseline (basal) versus stay in pit	OC during stay in pit lesser than basal condition; Subjects remained in pit till ambient O ₂ declined to 12% and CO ₂ rose to 7%; The maximum stay in pit for 18 hours when ambient CO ₂ was 7.7% and O ₂ 11.6%.	HR and BR rose when ambient CO ₂ reached to 5% in pit;
Craig Heller,	Yogi male (Proficient	Single practice	Stay in subterranean chamber for 4	Baseline (basal) versus	↓ of 40% in OC during the stay in	

Elsner et al. 1987 ⁽⁹⁹⁾	in <i>bhoogarbh smadhi-</i> (subterranean stay) (n=1);	on a single occasion	hours	stay in pit	chamber compared to basal baseline measured through gas volume meter	
Wallace 1970 ⁽¹²⁰⁾	Meditators with >6 months experience (n=15);	Single practice on a single occasion	Transcendental meditation (TM); 30 minutes meditation session	Baseline versus Meditation measured through	↓ of 20% in OC during meditation compared to baseline;	↑ of 103% in GSR ^{**} during onset of meditation;
Wallace, Benson et al. 1971 ⁽¹²¹⁾	Meditators with mean 29.4 months experience (n=36);	Single practice on a single occasion	Transcendental meditation; 30 minutes meditation session;	Baseline versus Meditation	↓ of 17% in OC (P<005) and ↓ of 15% in CO ₂ exhalation (P<005) during meditation;	↓ in HR and BR (Ps<05) during meditation; ↑ of 158% in GSR (P<005) during meditation;
Benson, Steinert et al. 1975 ⁽¹²²⁾	Meditators with >1 year experience (n=13);	Single practice on a single occasion	Transcendental meditation; 30 minutes meditation session;	Baseline versus Meditation	↓ of 5% in OC (P<001) and ↓ of 6% in CO ₂ exhalation (P<001) during meditation;	↓ in HR (P< 01) during meditation
Danucalov, Simões et al. 2008 ⁽¹³⁸⁾	Experienced yoga practitioner with >3 years' experience (n=9);	Multiple practices on a single occasion	Slow paced Pranayama with extended period of retention (Internal retention- I:R:E ⁺⁺ -1:4:2) and meditation; Each phase of 30 minutes;	Baseline versus Meditation and pranayama	↓ of 35% in OC (P<05) and ↓ 31.2% in CO ₂ exhalation (P<05) during meditation compared to baseline; ↓ of 49% in OC during meditation compared to <i>pranayama</i> ;	↓ of 8% in HR (P<05) during Meditation compared to baseline and <i>pranayama</i> ; ↑ of 20% in OC during <i>pranayama</i>

						compared to baseline;
Fenwick, Donaldson et al. 1977 ⁽¹³⁴⁾	Meditators with >22 months experience (n=11) and non-meditators (n=8);	Multiple practices on a single occasion	Transcendental meditation (TM) and listening music	Baseline versus meditation and listening to music in meditators and non-meditators; Comparison between groups	Non-significant drop in OC and CO ₂ exhalation during meditation in meditators and non-meditators both; Non-significant difference in reduction of OC between Meditation and listening music; Non-significant difference between groups;	No evidence of hypo-metabolism during Meditation in both the groups;
Warrenburg, Pagano et al. 1980 ⁽¹²³⁾	Regular meditators with mean 3.4 years' experience (n=9); Regular relaxation practitioners with mean 6.4 years' experience (n=9); Non practitioners (n=9);	Multiple practices on a single occasion	Transcendental meditation (TM); Progressive muscle relaxation (PMR); Non practitioner – listening music;	Control periods of closed eyes and reading book versus intervention; Comparison between groups	↓ in OC during TM 4%, during PMR 3.5%, in regular practitioners and 8.3% in non-practitioners (Ps<01) compared to periods of control; Non-significant difference between groups;	↓ HR during meditation/or relaxation (P<01);
Kesterson and Clinch 1989 ⁽¹²⁴⁾	Advanced meditators with mean 28 years experience (n=33); non-meditators	Multiple practices on a single occasion	Transcendental meditation (TM), Non-meditators - eyes closed relaxation	Baseline versus intervention; Comparison between the groups;	Similar significant drop in OC (P<002) during TM and relaxation; Non-significant difference between groups;	No traces of hypo-metabolism in either group;

Telles, Nagarathna et al. 1995 (117)	(n=10); Male meditators with >5 year experience (n=7);	Multiple practices on two occasions	<i>Aum</i> meditation session and sitting relaxed session; (Each session of 20 minutes on different days)	Baseline rest versus post meditation and eyes closed relaxation	Non-significant change in OC in post mediative session compared to baseline;	↓ in HR (P<001) during meditation
Vempati and Telles 1999 (101)	Male yoga practitioner with mean 23.9 months experience (n=40);	Multiple practices on two occasion	Yoga based Isometric relaxation and supine rest; (Each session of 10 minutes on different days)	Baseline rest versus post relaxation and supine rest	↓ of 23% (P<001) in OC in post yoga relaxation compared to baseline; ↓ of 7% in OC after supine rest;	↓ of 20.6% in BR (P<01) post yoga relaxation;
Vempati and Telles 2002 (100)	Male yoga practitioner with mean 30.2 months experience (n=35);	Multiple practices on two occasions	Yoga based guided relaxation session and supine rest session; (Each session of 10 minutes on different days)	Baseline rest versus post relaxation and supine rest	↓ of 25.2% in OC (P<001) in post yoga relaxation compared to baseline; ↓ of 7% in OC after supine rest;	↓ of 9.7% in HR (P<001), ↓ in LF ⁺⁺ (P<05), ↑ in HF ^{§§} (P<05) after relaxation compared to baseline
Ray, Pathak et al. 2011 (95)	Male yoga practitioners >6 years' experience (n=20);	Multiple practices on two occasions	<i>Hatha</i> Yoga session - comprising variety of yoga static postures interspersed with <i>Shavasana</i> , <i>pranayamas</i> and meditation practices; VO_{2max} session; (Each session on different days)	Rest sitting (<i>Sukhasana</i>) versus Meditation and <i>Aum</i> meditation	↓ of 15% in OC (P<05) during meditation and 4% during Om meditation compared to <i>Sukhasana</i> ;	↓ of 37% in BR (P<05) during Om Meditation ;

Throll 1982 (136)	Healthy non practitioner males (n=39)	15-weeks of regular practice multiple group	Meditation group– (n=21) Transcendental meditation , relaxation group – (n=18) Progressive muscle relaxation	Baseline versus immediately after practice of first session and then after 5, 10 and 15 weeks apart	Meditators displayed greater reduction in OC during the practice; reduction in OC more prominent in relaxation overtime of 15 weeks compared to meditation;	Reduction in HR in Meditation group prominent (P<05) compared to relaxation over time;
----------------------	---	--	---	---	--	--

Table 2: Summary of studies reporting changes in OC with mediation/relaxation practice(s)

Abbreviations used: – OC^{*} – oxygen consumption; HR[†] – heart rate; BPM[‡] - beats per minute; CO₂[§]- carbon dioxide; BMR^{||}- basal metabolic rate; BR[¶]- breath rate;

GSR^{**} -galvanic skin resistance; I:R:E^{††} – Inspiration:Retention:Expiration; LF^{‡‡} – low frequency; HF^{§§} – high frequency;

Table 3: Summary Of Studies Reporting Changes in Oxygen Consumption (OC*) With Asana, Integrated Practice(s)

Study Reference	Population	Study Design	Intervention	Comparators	Metabolic Measures	Cardio-respiratory and Other Measures
Rao 1962 (85)	Male yoga practitioners (n=6);	Single practice on a single occasion	Head stand posture	Baseline recumbent and standing erect versus head stand posture	↑ of 68% and 48% in OC during headstand posture compared to recumbent position and standing erect respectively;	
Rai and Ram 1993 (103)	Male yoga practitioners (n=10);	Single practice on a single occasion	<i>Virasana</i> (Hero Pose) (Subgroup 1 – with resting breath rate >10 breath/min (n=6); Subgroup 2- with resting breath rate <5 breath/min (n=4)	Baseline <i>Shavasana</i>	Subgroup 1: ↑ of 159% in OC (P<005) and ↑ of 223% in CO ₂ [†] exhalation (P<01) during <i>Virasana</i> in group with BR [‡] >10 breath/min; Subgroup 2: ↑ of 163% in OC (P<05) and ↑ of 166% in CO ₂ exhalation during <i>Viransana</i> in group with BR <5 breath/min:	↑ of 60% (P<0005) and 43% (P<001) in HR [§] in subgroup 1 and 2 respectively;
Rai, Ram et al. 1994 (102)	Male yoga practitioners (n=10);	Single practice on a single occasion	<i>Siddhasana</i> (Accomplished pose) (Subgroup 1 – with resting breath rate >10breath/min (n=6); Subgroup 2- with resting breath rate <5 breath/min (n=4)	Baseline <i>Shavasana</i>	Subgroup 1: ↑ of 27% in OC (P<01) and ↑ of 31% (P<.01) in CO ₂ exhalation during <i>Siddhasana</i> in group with >10 breath/min; Subgroup 2: ↑ of 21% in OC and ↑ of 25% in CO ₂ exhalation during <i>Siddhasana</i> in group with >5breath/min;	↑ of 13% (P<01) and of 15% (P<001) in HR in subgroup 1 and 2 respectively;

Sinha, Ray et al. 2004 (104)	Male yoga practitioners (n=21);	Single practice on a single occasion	Sun salutation (SS) – 12 dynamic postures preceded and followed by <i>Shavasana</i>	Comparison between each individual posture of SS and <i>Shavasana</i>	<p>↑ of 207% in OC during complete session of SS compared to <i>Shavasana</i>;</p> <p>↑ of 383% during 8th pose (Cobra) compared to <i>Shavasana</i>;</p> <p>OC higher (P<05) during backward bending poses (2nd, 4th, 5th and 8th poses) compared to forward bending poses (3rd and 11th);</p>	HR range - 83.5BPM ¹ to 101.6BPM during entire SS compared to 60.2BPM during <i>Shavasana</i> ;
Blank 2006 (125)	Female yoga practitioners (n=15);	Single practice on a single occasion	<i>Iyenger</i> Yoga posture sequences – warm-ups, 20 individual postures and releasing poses with <i>Shavasana</i>	Comparison between each individual posture and postures divided in sets (Back arch, inversion, standing, supine and seated) versus <i>Shavasana</i>	<p>↑ of OC during standing, back arc and inversion poses (P<05) compared to supine and seated posture;</p> <p>↑ of 300% of OC (P<05) during warrior pose III compared to <i>Shavasana</i>;</p> <p>↑ of 144% in OC during 65 minutes yoga session compared to <i>Shavasana</i></p>	Back arch poses 75% of HR _{max}
Hagins, Moore et al. 2007 (126)	Yoga practitioners with >1 year experience in yoga; 2 males; 28 females (n=20);	Multiple practices on a single occasion	<i>Ashtanga</i> yoga session of 56 minutes – Warm-up, sun salutation and non-sun salutation poses; Mild and moderate sub maximal exercise- treadmill walk at 2mph and	Baseline rest and mild to moderate exercise versus yoga session; Sun salutation versus non sun salutation poses	<p>↑ of 100% in OC (P<0001) during yoga session compared to rest.</p> <p>OC 14% lower during yoga sequence compared to mild exercise and 33% lower to</p>	↑ of 31% in HR (P<0001) during yoga compared to rest; Yoga sequence

			3mph;		moderate exercise (Ps<0001);	49.4% of HR _{max} ;
					OC 25% higher (P<001) during sun salutation compared to non-sun salutation poses;	HR 15% higher during sun salutation compared to non-sun salutation poses;
Telles, Reddy et al. 2000 (105)	Male yoga practitioner with >3 months experience (n=40);	Multiple practices on two occasions	Cyclic meditation-‘CM’ session and <i>Shavasana</i> session (Each session on different days)	Baseline rest versus post-practice session of CM and <i>Shavasana</i>	↓ of 32% in OC (P<001) post CM; ↓ of 10% in OC (P<05) post <i>Shavasana</i> compared to baseline	↓ of 28% (P<001) and 15% (P<05) in BR during post session in CM and <i>Shavasana</i> respectively
Sarang and Telles 2006 (106)	Male yoga practitioner with >3 months experience (n=50);	Multiple practices on two occasions	Cyclic meditation session – ‘CM’ (divided into 4 phases) and <i>Shavasana</i> session (Each session on different days);	Baseline rest versus CM session and <i>Shavasana</i> session; Baseline rest versus post practice session of CM and <i>Shavasana</i>	↑ of 31.3% in OC(P<001) during active phases of CM; ↓ of 19.4% in OC (P<001) post CM compared to baseline; ↓ of 7% in OC (P<001) post <i>Shavasana</i> ;	↑ up to 21% in BR (P<001) during CM and ↓ of 7% (P<05) in BR post CM compared to baseline;
DiCarlo, Sparling et al. 1995 (132)	Yoga practitioners with >1 year experience (n=10);	Multiple practices on two occasions	<i>Hatha</i> yoga - 12 standing postural sequence session; Sub maximal exercise session treadmill walk at 4 mph and VO2 _{max}	Sub maximal exercise and VO2 _{max} session versus <i>Hatha</i> yoga routine session	OC 26% lower during yoga sequences(P<05) compared to sub maximal exercise in first 8 th minute and remained lower during	HR 4% higher in during yoga sequence (P<05) compared to sub

			session; (Each session on different days);		complete session; Yoga session 34% of VO_{2max} and sub maximal exercise 46% of VO_{2max}	maximal exercise in 8 th minute and remained higher during complete yoga session;
Carroll, Blansit et al. 2003 (127)	Yoga practitioners with >3 months experience (n=13);	Multiple practices on two occasions	<i>Vinyasa</i> Yoga sequences and & VO_{2max}	VO_{2max} versus <i>Viniyasa</i> yoga	Yoga session >50% of VO_{2max}	Yoga session >77% of HR_{max} ;
Clay, Lloyd et al. 2005 (131)	Yoga practitioners with >1 month experience (n=30); 2 males, 28 females	Multiple practices on two occasions	<i>Hatha</i> yoga session – Warm-ups, sun salutation, non-sun salutation and cool down poses; Sub maximal exercise -treadmill walk at 3.5mph and VO_{2max} session; (Each session on different days)	Chair sitting, sub maximal exercise and VO_{2max} session versus <i>Hatha</i> yoga session	↑ of 114% in OC (P<05) during yoga session compared to chair sitting; OC 54% lower (P<05) during yoga session compared to sub maximal exercise; Yoga session 14.5% and sub maximal exercise 44.8% of VO_{2max} ; OC 82% higher (P<05) during sun salutation compared to non-sun salutation;	↑ of 24% in HR (P<06) during yoga session compared to chair sitting; HR 21% lower (P<05) during yoga session compared to sub maximal exercise; HR 20% higher (P<-5) during sun salutation compared to non-sun salutation;

Buranruk, La Grow et al. 2010 (137)	Middle aged non yoga practitioners (n=17);	Multiple practices on two occasions	Thai yoga session – warm-ups, sitting standing and lying poses; VO _{2max} ; (Each session on different days);	VO _{2max} session and baseline rest versus Thai yoga session calorimetry	↑ of 133% during Yoga session compared to rest; Yoga session 35.5% of VO _{2max} OC 46% higher (P<0001)during standing poses compared to sitting;	↑ of 16.6% in HR during Thai yoga compared to rest; HR during yoga sequence 50% of HR _{max} ; HR 10.6% higher (P<0001) during standing poses compared to sitting;
Ray, Pathak et al. 2011 (95)	Male yoga practitioners with >6 years' experience (n=20);	Multiple practices on two occasions	<i>Hatha</i> Yoga session - comprising variety of yoga static postures interspersed with <i>Shavasana</i> , <i>pranayamas</i> and meditation practices; VO _{2max} session; (Each session on different days)	VO _{2max} session and <i>Shavasana</i> versus individual yoga static postures;	↑ of 160% in OC (P<05)during plough pose-2 ↑ of 156% in OC (P<05) during bow pose compared to <i>Shavasana</i> ; Bow, Plough -1, plough -2 and shoulder stand pose 26.5% 25.9%, 24.6%, 22.7% respectively of VO _{2max} ; <i>Shavasana</i> 9.9% of Vo _{2max} ;	↑ of 108% in BR (P<05) during plough 1 & 2 compared to <i>Shavasana</i> ;

Table 3: Summary of studies reporting changes in OC with asana, integrated practice(s)

Abbreviations used: OC* - oxygen consumption; CO₂[†] - carbon dioxide; BR[‡] – breath rate; HR[§] – heart rate BPM^{||} - beats per minute;

Table 4: Summary of Studies Reporting Changes in Oxygen consumption (OC*) With Yoga and Physical Activity and Rest-

Study Reference	Population	Study design and duration	Intervention	Comparators	Metabolic Measures	Cardio-respiratory and Other Measures
Salgar, Bisen et al. 1975 ⁽¹⁰⁷⁾	Healthy males (n=38);	Multiple practices on single	6 months of regular lotus posture (n=10); resistance training (n=12); sedentary lifestyle (n=16);	Comparison between groups during mild and moderate level ergometer exercise	At mild level exercise the OC was lowest in lotus group followed by exercisers and non-exercisers; At moderate level exercise OC lowest in exercisers followed by lotus and non-exerciser group	
Bhatnagar, Ganguly et al. 1978 ⁽¹⁰⁸⁾	Healthy non yoga practitioners (n=20);	6-month Cohort study of multiple practices	Regular integrated yoga practices	Pre yoga intervention sub maximal exercise versus fixed intensity sub maximal exercise after 1,3 and 6 months of yoga practice	Progressive increase in OC during sub maximal exercise (P< 05) in 3 and 6 months compared to pre intervention;	↓ in resting body core temperature (P<05, P<001,P<001) in 1, 3 and 6 months compared to pre intervention;
Joseph, Sridharan et al. 1981 ⁽¹⁰⁹⁾	Healthy male non yoga practitioners (n=10);	3-month Cohort study of multiple practices	Regular integrated yoga practices	Pre yoga intervention rest versus post yoga intervention rest	Non- significant decrease in resting OC;	↓ of 7.8% in HR [†] (P<001); ↓ in SBP [‡] /DBP [§] (P<01); ↓ in blood

						glucose (P<05) and blood cholesterol (P<01);
Raju, Kumar et al. 1986 ⁽¹¹⁰⁾	Non yoga practitioners (n=12);	3-month Cohort study of multiple practices	Regular integrated yoga practice	Pre yoga intervention rest versus post yoga intervention rest versus; Pre yoga intervention sub maximal exercise versus Sub maximal exercise after 20 days and 3 months of yoga practice	Non-significant change in resting OC in either gender; ↓ of 41% in OC (P<05) during submaximal exercise after 20 days and ↓ of 36% in OC (P<05) during sub-maximal exercise after 3 months in males only;	↓ of 65.5% in blood lactate (P<05) in males at same exercise workload after 3 months; No significant changes in females;
Balasubramanian and Pansare 1991 ⁽¹¹⁶⁾	Healthy non yoga practitioners (n=17);	6-week Cohort study of multiple practices	Integrated yoga practice	Pre yoga intervention VO _{2max} versus post yoga intervention VO _{2max}	↑ of 17% in VO _{2max} (P<005);	
Raju, Prasad et al. 1997 ⁽¹¹¹⁾	Healthy female non yoga practitioner (n=6);	4-week Cohort study of multiple practices	Integrated yoga practice	Pre yoga intervention VO _{2max} versus exercise versus post yoga intervention VO _{2max}	↓ of 14% in OC (P<05) per unit of work load; ↑ of 21% in maximal work load (P<05) in post yoga intervention compared to pre yoga intervention	↓ of 6% in HR (P<05) post yoga intervention; ↓ in body fat & weight (P<05);
Tran, Holly et	Healthy non yoga	8-week	Integrated yoga session	Pre yoga intervention	↑ of 10% in VO _{2max} (P<01);	↑ in muscular

al. 2001 (128)	practitioners (n =10);	Cohort study of multiple practices		VO _{2max} versus post yoga intervention VO _{2max}		strength (P<05), muscular endurance (P< 01) and flexibility (P<001);
Ramos- Jiménez, Hernández- Torres et al. 2011 (142)	Female middle and old aged yoga practitioners with >3 years' experience (n=13);	11-week Cohort study of multiple practices	Integrated intensive yoga training; (middle aged practitioners with mean age 43.2 years = 4; older aged practitioners with mean age 62.2 years =9);	Pre yoga intervention VO _{2max} versus post yoga intervention VO _{2max}	↑ of 3% in VO _{2max} (P<05) in middle aged group and ↑ of 17% in VO _{2max} (P<05) in older aged group;	Increase in HR _{max} (P<05) in both groups; Improvement in Lipid profile and blood glucose and BMI (Ps<05) in both groups;
Raju, Madhavi et al. 1994 (112)	Healthy male non- yoga practitioners (n=28);	24-month NRCT of multiple practices	Yoga group-Pranayama and <i>Shavasana</i> along with regular sports workouts (n=14) Control – Regular sports workouts (n=14) (Each group further sub grouped in Phase 1 and 2 of submaximal (n=12) and maximal exercise (n=16) of duration 12 months and 24 months respectively)	Pre intervention rest versus post yoga intervention rest (phase 1&2); Pre intervention sub maximal and maximal exercise versus post yoga intervention sub maximal and maximal exercise	Phase 1 - ↓ of 38% in OC at resting state in yoga group; ↓ of 51% in OC (P<05) per unit work load with sub maximal exercise in yoga group after intervention; No change in controls either in rest or during exercise; Phase 2- No change in resting OC in either yoga or control group; ↓ of 34% in OC (P<05) per unit	Phase 1 - ↓ of 49% in resting blood lactate (P<01) in yoga group; Phase 2 - ↓ of 37% in resting blood lactate (P<05) in yoga group; ↓ of 61% in

					work load with maximal exercise in yoga group after intervention; No change in control;	exercise blood lactate after 24 months compared to pre intervention in yoga group;
U. S. Ray, Sinha, Tomer, Pathak, & et al., 2001 (115)	Healthy male non-yoga practitioners (n=28);	6-month RCT** of multiple practices	Yoga group- Integrated yoga practises (n=17); Physical training as per army program(n=11);	Pre intervention VO2 _{max} versus post intervention VO2 _{max}	↑ of 6.7% in VO2 _{max} (P<05) in yoga group; No change in physical training group;	↓ in body fat and body weight (Ps<01) in yoga group;
Nayar, Mathur et al. 1975 (113)	Healthy male non-yoga practitioners (n=53);	12-month RCT of multiple practices	Yoga group- Integrated yoga with regular physical training (n=18); Athletic group- Athletics with regular physical training (n=17); Control – regular physical training (n=18);	Pre intervention rest versus post intervention rest: Pre intervention sub-maximal versus post intervention sub maximal exercise	Non- significant change in OC at rest in either group; Non- significant changes in OC during sub-maximal exercise in either group;	↑ of 29% in vital capacity (P<01) and 5% in FEV1 ^{††} (P<05) in yoga group; ↑ of 46% in breath-hold time (P<01) in yoga group;
Selvamurthy, Ray et al. 1988 (114)	Healthy male non-yoga practitioners (n=30);	6-month RCT of multiple practices	Yoga group- Integrated training (n=15); Physical training 'PT' group- running, games, flexibility and pull-ups (n=15);	Pre yoga intervention sub maximal exercise versus post yoga intervention sub maximal exercise	↓ of 5.7% in OC (P<05) in yoga group; Non -significant change in PT group;	↓ of 7% in HR (P<01) in yoga group;

Bowman, Clayton et al. 1997 (133)	Sedentary healthy elderly subjects > 62 years (n=40);	6-week RCT of multiple practices	Yoga group- Integrate yoga (n=20); Aerobic group- Bicycle based aerobic training (n=20);	Pre yoga intervention sub maximal exercise versus post yoga intervention sub maximal exercise	↑ of 13% in VO _{2max} (P<01) in yoga group and ↑ of 24% in VO _{2max} (P<01) in aerobic group;	↓ of 11.6% in HR (P<05) in yoga group; No change in aerobic group; ↑ in baroreflex sensitivity (P<01) in yoga group; No significant change in HRV ^{##} in either groups;
Pullen, Nagamia et al. 2008 (130)	Patients with congestive heart failure 'CHF' (n=19);	8-week RCT of multiple practices	Yoga group- Integrated yoga practices along with standard medical therapy (n=9); Control – Standard medical therapy with general awareness (n=10);	Pre intervention VO _{2max} versus post intervention VO _{2max}	↑ of 17% in VO _{2max} (P<02) in yoga group; No change in controls;	Improvement of 25.7% in quality of life scores (P<.005) in yoga group
Tracy and Hart 2012 (129)	Sedentary healthy non yoga practitioners (n=21);	8-week RCT of multiple practices	<i>Bikram</i> yoga- 26 series of postures in heated 35 ⁰ -40 ⁰ C humidified studio (n=10); Waitlist control (n=11)	Pre yoga intervention VO _{2max} versus post yoga intervention VO _{2max}	No change VO _{2max} after yoga training;	↑ of 23.8% in sit and reach score (P<001) and shoulder flexibility (P<05) with yoga;
Chaya, Kurpad et al. 2006 (70)	Non yoga (NY) and regular yoga practitioners (YP)	Multiple practices on a single	YP - Regular integrated yoga practice (n=55); NY - (n=49)	Yoga practitioners versus Non yoga practitioners at rest (basal state)	Basal OC - 19.3% less in female YPs and 10.7% less in male YPs (Ps<001) compared to NYs;	BR ^{¶¶} 19.6% less (P<001) in female YPs and 19%

	with >6 months experience n=104);	occasion			Basal CO ₂ ⁵⁵ - 12.7% less in female YPs and 14.3% less in male YPs (Ps<05) compared to NYs; BMR ^{III} in YP 15% less (P<001) in YP compared to NY and 13% less than the predicted by WHO/FAO/UNU;	(P<001) in male in YPs compared to NYs;
Chaya and Nagendra 2008 ⁽⁷¹⁾	Non-yoga (NY) and regular yoga practitioners (YP) with >6 months experience (n=88);	Multiple practices on a single occasion	YP - Regular integrated yoga practice (n= 51); NY – (n=37)	Yoga practitioners versus non yoga practitioners at rest at 6am (basal) and 9 pm (pre sleep)	Basal OC - 22% less (P<005) in female YPs and 10.7% less (P<05) in male YPs compared to NY females and males respectively; Pre-sleep OC - 17% less in female YPs and 6.7% in male YPs (non-significant) compared to NY females and males respectively; Basal CO ₂ – 15.3% less in female YPs and 14.8% less in male YPs (Ps<05) compared to NYs; Pre sleep CO ₂ - 13.2% less in female YPs and 8.3% in male YPs compared to NYs	BR 23.3% less (P<005) in female YPs and 15.6% less (P<05) in male YPs during morning compared to NYs;

Table 4: Summary of studies reporting changes in OC with yoga and physical; activity

Abbreviations used - OC* - oxygen consumption; HR[†] – heart rate; SBP[‡]- systolic blood pressure; DBP[§] - diastolic blood pressure; BMI^{||} –body mass index, NRCT[¶] – non randomised controlled trials; RCT^{**} – randomised controlled trials; FEV1^{††} – forced expiratory volume in 1 second; HRV^{‡‡} – heart rate variability; CO₂⁵⁵- carbon dioxide exhalation; BMR^{III}- basal metabolic rate; BR^{¶¶} - breath rate

References

1. Haugen HA, Chan L-N, Li F. Indirect Calorimetry: A Practical Guide for Clinicians. *Nutr Clin Pract*. 2007 August 1, 2007;22(4):377-88.
2. Bonnet MH, Arand DL. Insomnia, metabolic rate and sleep restoration. *J Intern Med*. 2003;254(1):23-31.
3. McArdle WD, Katch FI, Katch VL. Exercise physiology : nutrition, energy, and human performance. Philadelphia [u.a.]: Wolters Kluwer, Lippincott Williams & Wilkins; 2010.
4. Olshansky SJ, Rattan S. What determines longevity: metabolic rate or stability. *Discovery Medicine*. 2005;5(28):359-62.
5. Epel ES. Psychological and metabolic stress: A recipe for accelerated cellular aging? *Hormones*. 2009;8(1):7-22.
6. Levine JA. Measurement of energy expenditure. *Public Health Nutr*. 2005;8(7 A):1123-32.
7. Glass S, Dwyer GB, American College of Sports M. ACSM'S metabolic calculations handbook. Philadelphia, Pa.: Lippincott Williams & Wilkins; 2007.
8. Jumpertz R, Hanson RL, Sievers ML, Bennett PH, Nelson RG, Krakoff J. Higher Energy Expenditure in Humans Predicts Natural Mortality. *J Clin Endocrinol Metab*. 2011 March 30, 2011.
9. Ruggiero C, Metter EJ, Melenovsky V, Cherubini A, Najjar SS, Ble A, et al. High Basal Metabolic Rate Is a Risk Factor for Mortality: The Baltimore Longitudinal Study of Aging. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2008 July 1, 2008;63(7):698-706.
10. Carroll D, Phillips AC, Balanos GM. Metabolically exaggerated cardiac reactions to acute psychological stress revisited. *Psychophysiology*. 2009;46(2):270-5.
11. Balanos GM, Phillips AC, Frenneaux MP, McIntyre D, Lykidis C, Griffin HS, et al. Metabolically exaggerated cardiac reactions to acute psychological stress: The effects of resting blood pressure status and possible underlying mechanisms. *Biol Psychiatry*. 2010;85(1):104-11.
12. Carroll D, Rick Turner J, Hellawell JC. Heart Rate and Oxygen Consumption during Active Psychological Challenge: The Effects of Level of Difficulty. *Psychophysiology*. 1986;23(2):174-81.
13. Turner JR, Carroll D. Heart Rate and Oxygen Consumption during Mental Arithmetic, a Video Game, and Graded Exercise: Further Evidence of Metabolically-Exaggerated Cardiac Adjustments? *Psychophysiology*. 1985;22(3):261-7.
14. Schmidt WD, O'Connor PJ, Cochrane JB, Cantwell M. Resting metabolic rate is influenced by anxiety in college men. *J Appl Physiol*. 1996;80(2):638-42.
15. Poehlman ET, Scheffers J, Gottlieb SS, Fisher ML, Vaitekevicius P. Increased Resting Metabolic Rate in Patients with Congestive Heart Failure. *Ann Intern Med*. 1994 December 1, 1994;121(11):860-2.
16. Bernardi M, Macaluso A, Sproviero E, Castellano V, Coratella D, Felici F, et al. Cost of walking and locomotor impairment. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology*. 1999 Apr;9(2):149-57. PubMed PMID: 10098715. eng.
17. Hommes MJ, Romijn JA, Endert E, Sauerwein HP. Resting energy expenditure and substrate oxidation in human immunodeficiency virus (HIV)-infected asymptomatic men: HIV affects host metabolism in the early asymptomatic stage. *Am J Clin Nutr*. 1991 August 1, 1991;54(2):311-5.
18. Wouters EFM. Nutrition and Metabolism in COPD*. *CHEST Journal*. 2000;117(5_suppl_1):274S-80S.
19. Poehlman ET, Melby CL, Badylak SF, Calles J. Aerobic fitness and resting energy expenditure in young adult males. *Metabolism*. 1989;38(1):85-90.
20. Tarantino G, Marra M, Contaldo F, Pasanisi F. Basal metabolic rate in morbidly obese patients with non-alcoholic fatty liver disease. *Clinical and Investigative Medicine*. 2008;31(1):E24-E9.
21. KRESS J, POHLMAN A, ALVERDY J, HALL J. The Impact of Morbid Obesity on Oxygen Cost of Breathing (\dot{V}_O 2RESP) at Rest. *Am J Respir Crit Care Med*. 1999 September 1, 1999;160(3):883-6.
22. Ravussin E, Burnand B, Schutz Y, Jéquier E. Twenty-four-hour energy expenditure and resting metabolic rate in obese, moderately obese, and control subjects. *Am J Clin Nutr*. 1982 March 1, 1982;35(3):566-73.
23. Regensteiner JG, Sippel J, McFarling ET, Wolfel EE, Hiatt WR. Effects of non-insulin-dependent diabetes on oxygen consumption during treadmill exercise. *Med Sci Sports Exerc*. 1995;27(5):661-7.
24. Huang K-C, Kormas N, Steinbeck K, Loughnan G, Caterson ID. Resting Metabolic Rate in Severely Obese Diabetic and Nondiabetic Subjects[ast][ast]. *Obesity*. 2004 05//print;12(5):840-5.

25. Fisher P, Kleinerman JI. TOTAL OXYGEN CONSUMPTION AND METABOLIC RATE OF PATIENTS IN DIABETIC ACIDOSIS 1. *The Journal of Clinical Investigation*. 1952;31(1):126-30.
26. Horstmann P. The Oxygen Consumption in Diabetes Mellitus. *Acta Med Scand*. 1951;139(4):326-30.
27. Snodgrass JJ, Leonard WR, Sorensen MV, Tarskaia LA, Mosher MJ. The influence of basal metabolic rate on blood pressure among indigenous Siberians. *Am J Phys Anthropol*. 2008 Oct;137(2):145-55. PubMed PMID: 18470897. Epub 2008/05/13. eng.
28. Kunz I, Schorr U, Klaus S, Sharma AM. Resting Metabolic Rate and Substrate Use in Obesity Hypertension. *Hypertension*. 2000 July 1, 2000;36(1):26-32.
29. Rosenkrantz Ja MC. Basal metabolic rate in hypertensive vascular disease. *Arch Intern Med*. 1947;80(1):81-8.
30. Vaccarino V, Bremner JD. Stress response and the metabolic syndrome. *Cardiology*. 2005;11(Part 2):1.
31. Licht CMM, Vreeburg SA, van Reedt Dortland AKB, Giltay EJ, Hoogendijk WJG, DeRijk RH, et al. Increased sympathetic and decreased parasympathetic activity rather than changes in hypothalamic-pituitary-adrenal axis activity is associated with metabolic abnormalities. *J Clin Endocrinol Metab*. 2010;95(5):2458-66.
32. Lambert EA, Lambert GW. Stress and Its Role in Sympathetic Nervous System Activation in Hypertension and the Metabolic Syndrome. *Curr Hypertens Report*. 2011:1-5.
33. Kyrou I, Tsigos C. Stress hormones: physiological stress and regulation of metabolism. *Current Opinion in Pharmacology*. 2009;9(6):787-93.
34. Benson H. *The relaxation response*. New York: Morrow; 1975.
35. Dusek JA, Benson H. Mind-body medicine: a model of the comparative clinical impact of the acute stress and relaxation responses. *Minn Med*. 2009 May;92(5):47-50. PubMed PMID: 19552264. Pubmed Central PMCID: PMC2724877. Epub 2009/06/26. eng.
36. Storey KB, Storey JM. Metabolic rate depression and biochemical adaptation in anaerobiosis, hibernation and estivation. *Q Rev Biol*. 1990:145-74.
37. Wolsko Pea. Use of mind-body medical therapies. *J Gen Intern Med*. 2004;19(1):43-50.
38. Young JDE, Taylor E. Meditation as a voluntary hypometabolic state of biological estivation. *News Physiol Sci*. 1998;13(3):149-53.
39. Dillbeck MC, Orme-Johnson DW. Physiological differences between transcendental meditation and rest. *Am Psychol*. 1987;42(9):879-81. PubMed PMID: 614372829; 1988-03610-001. English.
40. Delmonte MM. Physiological responses during meditation and rest. *Biofeedback Self Regul*. 1984;9(2):181-200.
41. Lehrer P, Schocket S, Carrington P, et al. Psychophysiological and cognitive responses to stressful stimuli in subjects practicing progressive relaxation and clinically standardized meditation. *Behav Res Ther*. 1980;18(4):293-303.
42. Elson BD, Hauri P, Cunis D. Physiological changes in yoga meditation. *Psychophysiology*. 1977;14(1):52-7.
43. Benson H, Dryer T, Hartley LH. Decreased VO₂ consumption during exercise with elicitation of the relaxation response. *J Human Stress*. 1978 Jun;4(2):38-42. PubMed PMID: 351050. Epub 1978/06/01. eng.
44. Telles S, Patra S, Montesoo S, Naveen KV. Effect of yoga on somatic indicators of stress in healthy volunteers. *J Indian Psychol*. 2008;26(1-2):52-7.
45. Parshad O. Role of yoga in stress management. *West Indian Med J*. 2004;53(3):191-4.
46. Sanghani S DA, Herring P, et al. A Pilot Study: Can a Short-term Complementary and Alternative Medicine Intervention Combat Stress? *Californian J Health Promot*. 2008;6(2):73-8.
47. Hartfiel N HJ, Khalsa SB, et al. The effectiveness of yoga for the improvement of well-being and resilience to stress in the workplace. *Scand J Work Environ Health*. 2011;37(1):70.
48. Gopal A, Mondal S, et al. . Effect of Integrated Yoga Practice on immune response in examination stress- A preliminary Study. *International journal of yoga*. 2011;4(1):26-32.
49. Malathi A, Damodaran A. Stress due to exams in medical students - Role of yoga. *Indian J Physiol Pharmacol*. 1999;43(2):218-24.
50. DiNardo MM. Mind-body therapies in diabetes management. *Diabetes Spectr*. 2009;22(1):30-4.
51. Gupta N, Khera S, Vempati RP, et al. Effect of yoga based lifestyle intervention on state and trait anxiety. *Indian J Physiol Pharmacol*. 2006;50(1):41-7.

52. Agte VV, Chiplonkar SA. Sudarshan Kriya Yoga for Improving Antioxidant Status and Reducing Anxiety in Adults. *Alter Comp Therp.* 2008;14(2):96-100.
53. Javnbakht M, Hejazi Kenari R, Ghasemi M. Effects of yoga on depression and anxiety of women. *Complement Ther Clin Pract.* 2009;15(2):102-4.
54. Vedamurthachar A, Janakiramaiah N, Hegde JM, et al. Antidepressant efficacy and hormonal effects of Sudarshana Kriya Yoga (SKY) in alcohol dependent individuals. *J Affect Disord.* 2006;94(1-3):249-53.
55. Yoshihara K, Hiramoto T, Sudo N, et al. Profile of mood states and stress-related biochemical indices in long-term yoga practitioners. *BioPsychoSocial Medicine.* 2011;5.
56. Wood C. Mood change and perceptions of vitality: A comparison of the effects of relaxation, visualization and yoga. *J R Soc Med.* 1993;86(5):254-8.
57. Aljasir B, Bryson M, Al-Shehri B. Yoga practice for the management of type II diabetes mellitus in adults: A systematic review. *Evid Based Complement Alternat Med.* 2010;7(4):399-408.
58. Matsuda K, Ueno S, Oura N, et al. Yoga Effects on Moods among Women with Infants. *J Psychosom Obstet Gynaecol.* 2007;28:77.
59. Carter J, Byrne G. A two year study of the use of yoga in a series of pilot studies as an adjunct to ordinary psychiatric treatment in a group of Vietnam War veterans suffering from post traumatic stress disorder. Online document at: www.Therapywithyoga.com Accessed November. 2004;27.
60. Rosenthal JZ, Grosswald S, Ross R, et al. Effects of transcendental meditation in veterans of Operation Enduring Freedom and Operation Iraqi Freedom with posttraumatic stress disorder: a pilot study. *Mil Med.* 2011 Jun;176(6):626-30. PubMed PMID: 21702378. eng.
61. Brooks JS, Scarano T. Transcendental Meditation in the Treatment of Post-Vietnam Adjustment. *J Couns Dev.* 1985;64(3):212-5.
62. Telles S, Naveen K, Dash M. Yoga reduces symptoms of distress in tsunami survivors in the andaman islands. *Evid Based Complement Alternat Med.* 2007;4(4):503-10.
63. Descilo T, Vedamurtachar A, Gerbarg PL, et al. Effects of a yoga breath intervention alone and in combination with an exposure therapy for post-traumatic stress disorder and depression in survivors of the 2004 South-East Asia tsunami [NCT00290225]. *Acta Psychiatr Scand*2010. p. 289-300.
64. Gerbarg PL, Brown RP. Yoga: A breath of relief for Hurricane Katrina refugees. *Current Psychiatry.* 2005;4(10):55-67.
65. Telles S, Singh N, Joshi M, Balkrishna A. Post traumatic stress symptoms and heart rate variability in Bihar flood survivors following yoga: A randomized controlled study. *BMC psychiatry.* 2010;10.
66. Li AW, Goldsmith CAW. The effects of yoga on anxiety and stress. *Altern Med Rev.* 2012;17(1):21-35.
67. Chong CS, Tsunaka M, Tsang HW, Chan EP, Cheung WM. Effects of yoga on stress management in healthy adults: A systematic review. *Altern Ther Health Med.* 2011 //;17(1):32-8.
68. Bharshankar JR, Bharshankar RN, Deshpande VN, Kaore SB, Gosavi GB. Effect of yoga on cardiovascular system in subjects above 40 years. *Indian J Physiol Pharmacol.* 2003;47(2):202-6.
69. Gopal KS, Bhatnagar OP, Subramanian N, Nishith SD. Effect of yogasanas and pranayamas on blood pressure, pulse rate and some respiratory functions. *Indian J Physiol Pharmacol.* 1973;17(3):273-6.
70. Chaya M, Kurpad A, Nagendra H, Nagarathna R. The effect of long term combined yoga practice on the basal metabolic rate of healthy adults. *BMC Complement Altern Med.* 2006;6(1):28. PubMed PMID: doi:10.1186/1472-6882-6-28.
71. Chaya M, Nagendra H. Long-term effect of yogic practices on diurnal metabolic rates of healthy subjects. *International journal of yoga.* 2008;1(1):27.
72. Balk JL. Yoga for weight loss. *Altern Med Alert.* 2011;14(5):49-53.
73. Seo DY, Lee S, Figueroa A, et al. Yoga training improves metabolic parameters in obese boys. *Korean J Physiol Pharmacol.* 2012;16(3):175-80.
74. Vyas R, Raval KV, Dikshit N. Effect of Raja yoga meditation on the lipid profile of post-menopausal women. *Indian J Physiol Pharmacol.* 2008 Oct-Dec;52(4):420-4. PubMed PMID: 19585761. Epub 2009/07/10. eng.
75. Telles S, Visweswaraiyah NK, Balkrishna A. Serum leptin, cholesterol and blood glucose levels in diabetics following a yoga and diet change program. *Med Sci Monit.* 2010;16(3):LE4-LE5.
76. Pal A, Srivastava N, Tiwari S, et al. Effect of yogic practices on lipid profile and body fat composition in patients of coronary artery disease. *Complement Ther Med.* 2011;19(3):122-7.

77. Sahay BK. Role of yoga in diabetes. *J Assoc Physicians India*. 2007;55(FEB.):121-6.
78. Mohta N, Agrawal RP, Kochar DK, Kothari RP, Sharma A. Influence of Yogic Treatment on Quality of Life Outcomes, Glycemic Control, and Risk Factors in Diabetes Mellitus: Randomized Controlled Trial. *Explore (NY)*. 2009;5(3):147.
79. Bhavanani AB, Sanjay ZZ, Madanmohan. Immediate effect of sukha pranayama on cardiovascular variables in patients of hypertension. *Int J Yoga Therap*. 2011 (21):73-6. PubMed PMID: 927688817; 22398346. eng.
80. Okonta NR. Does yoga therapy reduce blood pressure in patients with hypertension?: an integrative review. *Holist Nurs Pract*. 2012 May-Jun;26(3):137-41. PubMed PMID: 22517349. eng.
81. Shapiro D, Surwit RS. Operant conditioning: a new theoretical approach in psychosomatic medicine. *Int J Psychiatry Med*. 1974 Fall;5(4):377-87. PubMed PMID: 4618839. Epub 1974/01/01. eng.
82. Cohen BE, Chang AA, Grady D, et al. Restorative yoga in adults with metabolic syndrome: A randomized, controlled pilot trial. *Metabolic syndrome and related disorders*. 2008;6(3):223-9.
83. Khatri D, Mathur KC, Gahlot S, Jain S, Agrawal RP. Effects of yoga and meditation on clinical and biochemical parameters of metabolic syndrome. *Diabetes Res Clin Pract*. 2007;78(3):e9-e10.
84. Lee JA, Kim JW, Kim DY. Effects of yoga exercise on serum adiponectin and metabolic syndrome factors in obese postmenopausal women. *Menopause*. 2012;19(3):296-301.
85. Rao S. The Metabolic Cost of Head-Stand Posture. *J Appl Physiol*. 1962;17(1):117 - 8.
86. Rao S. Oxygen consumption during yoga-type breathing at altitudes of 520m. and 3,800m. *Indian J Med Res*. 1968 May;56(5):701-5. PubMed PMID: 5697820. Epub 1968/05/01. eng.
87. Karambelkar PV, Deshapande, R.R., & Bhole, M.V. Some respiratory studies in respect of kapalbhati and voluntary hyperventilation. *Yoga Mimamsa*. 1982;21(1&2):54-8.
88. Karambelkar PV, Deshapande, R.R., & Bhole, M.V. Some Respiratory Studies on Bhastrika Pranayama with Internal and External Retention. *yoga Mimamasa*. 1982;21(3 & 4):14-20.
89. Karambelkar PV, & Bhole, M.V. Respiratory studies during kapalbhati for 1, 2, 3 and 5 minutes. *Yoga Mimamsa*. 1988;27((1&2)):69-74.
90. Karambelkar PV, Deshapande, R.R., & Bhole, M.V. Some respiratory studies of ujjayi and bhastrika pranayama with bahya kumbhak. *Yoga Mimamasa*. 1983;22(3 & 4):7-12.
91. Karambelkar PV, Deshpande, R.R., Bhole, M.V. Oxygen Consumption during Ujjayi Pranayama. *Yoga Mimamsa*. 1983;21(3&4):7-13.
92. Telles S, Desiraju T. Oxygen consumption during pranayamic type of very slow-rate breathing. *Indian J Med Res*1991. p. 357-63.
93. Telles S, Nagarathna R, Nagendra HR. Physiological measures of right nostril breathing. *J Altern Complement Med*. 1996 Winter;2(4):479-84. PubMed PMID: 9395677. Epub 1996/01/01. eng.
94. Prasad KVV, Venkata Ramana Y, Raju PS, et al. Energy cost and physiological efficiency in male yoga practitioners. *J Exerc Physiol Online*. 2001;4(3):38-44.
95. Ray US, Pathak A, Tomer OS. Hatha Yoga practices: Energy expenditure, respiratory changes and intensity of exercise. *Evid Based Complement Alternat Med*. 2011;2011.
96. Telles S, Nagarathna R, Nagendra HR. Breathing through a particular nostril can alter metabolism and autonomic activities. *Indian J Physiol Pharmacol*. 1994;38(2):133-7.
97. Anand B, Chhina G, Singh B. Studies on Shri Ramanand Yogi during his stay in an air-tight box. *Indian J Med Res*. 1961.
98. Karambelkar PV, Vinekar SL, Bhole MV. Studies on human subjects staying on an air-tight pit. *Indian J Med Res*. 1968 Aug;56(8):1282-8. PubMed PMID: 5711607. Epub 1968/08/01. eng.
99. Craig Heller H, Elsner R, Rao N. Voluntary hypometabolism in an indian yogi. *J Therm Biol*. 1987;12(2):171-3.
100. Vempati R, Telles S. Yoga based guided relaxation reduces sympathetic activity in subjects based on baseline levels. *Psychol Rep*. 2002;90:487-94.
101. Vempati R, Telles S. Yoga based isometric relaxation versus supine rest: A study of oxygen consumption, breath rate and volume and autonomic measures. *J Indian Psychol*. 1999;17(2):46-52.
102. Rai L, Ram K, Kant U, Madan SK, Sharma SK. Energy expenditure and ventilatory responses during Siddhasana - A yogic seated posture. *Indian J Physiol Pharmacol*. 1994;38(1):29-33.

103. Rai L, Ram K. Energy expenditure and ventilatory responses during Virasana--a yogic standing posture. *Indian J Physiol Pharmacol.* 1993;37(1):45-50.
104. Sinha B, Ray U, Pathak A, et al. Energy cost and cardiorespiratory changes during the practice of Surya Namaskar. *Indian J Physiol Pharmacol.* 2004;48(2):184 - 90.
105. Telles S, Reddy S, Nagendra H. Oxygen consumption and respiration following two yoga relaxation techniques. *Appl Psychophysiol Biofeedback.* 2000;25(4):221-7.
106. Sarang PS, Telles S. Oxygen consumption and respiration during and after two yoga relaxation techniques. *Appl Psychophysiol Biofeedback.* 2006;31(2):143-53.
107. Salgar DC, Bisen VS, Jinturkar MJ. Effect of padmasana. A yogic exercise on muscular efficiency. *Indian J Med Res.* 1975;63(6):768-72.
108. Bhatnagar OP, Ganguly AK, Anantharaman V. Influence of Yoga training on thermoregulation. *Indian J Med Res.* 1978;67(5):844-7.
109. Joseph S, Sridharan K, Patil SKB. Study of some physiological and biochemical parameters in subjects undergoing yogic training. *Indian J Med Res.* 1981;74(1):120-4.
110. Raju PS, Kumar KA, Reddy SS, et al. Effect of yoga on exercise tolerance in normal healthy volunteers. *Indian J Physiol Pharmacol.* 1986;30(2):121-32.
111. Raju PS, Prasad KVV, Venkata RY, et al. Influence of intensive yoga training on physiological changes in 6 adult women: A case report. *J Altern Complement Med.* 1997;3(3):291-5.
112. Raju PS, Madhavi S, Prasad KV, et al. Comparison of effects of yoga & physical exercise in athletes. *Indian J Med Res.* 1994;81-6. PubMed PMID: CN-00104974.
113. Nayar HS, Mathur RM, Sampath Kumar R. Effects of yogic exercises on human physical efficiency. *Indian J Med Res.* 1975;63(10):1369-76.
114. Selvamurthy W, Ray US, Hegde KSea. Physiological responses to cold (10° C) in men after six months' practice of yoga exercises. *Int J Biometeorol.* 1988;32(3):188-93.
115. Ray US, Sinha B, Tomer OS, Pathak A, et al. Aerobic capacity & perceived exertion after practice of Hatha yogic exercises. *Indian J Med Res.* 2001;114:215.
116. Balasubramanian B, Pansare MS. Effect of yoga on aerobic and anaerobic power of muscles. *Indian J Physiol Pharmacol.* 1991;35(4):281-2.
117. Telles S, Nagarathna R, Nagendra HR. Autonomic changes during "OM" meditation. *Indian J Physiol Pharmacol.* 1995 Oct;39(4):418-20. PubMed PMID: 8582759. Epub 1995/10/01. eng.
118. Miles WR, Behanan KT. A metabolic study of three unusual learned breathing patterns practiced in the cult of Yoga. *Am J Psychol.* 1934;109:74-5.
119. Miles W. Oxygen consumption during three yoga-type breathing patterns. *J Appl Physiol.* 1964;19(1):75.
120. Wallace RK. Physiological effects of transcendental meditation. *Science.* 1970;167(3926):1751-4.
121. Wallace RK, Benson H, Wilson AF. A wakeful hypometabolic physiologic state. *Am J Psychol.* 1971;221(3):795-9.
122. Benson H, Steinert RF, Greenwood MM, et al. Continuous measurement of O₂ consumption and CO₂ elimination during a wakeful hypometabolic state. *J Human Stress.* 1975;1(1):37-44.
123. Warrenburg S, Pagano RR, Woods M, et al. A comparison of somatic relaxation and EEG activity in classical progressive relaxation and transcendental meditation. *J Behav Med.* 1980;3(1):73-93.
124. Kesterson J, Clinch NF. Metabolic rate, respiratory exchange ratio, and apneas during meditation. *Am J Physiol Regul Integr Comp Physiol.* 1989 March 1, 1989;256(3):R632-R8.
125. Blank SE. Physiological responses to iyengar yoga performed by trained practitioners. *J Exerc Physiol Online.* 2006;9(1):7-23.
126. Hagins M, Moore W, Rundle A. Does practicing hatha yoga satisfy recommendations for intensity of physical activity which improves and maintains health and cardiovascular fitness? *BMC Complement Altern Med.* 2007;7.
127. Carroll J, Blansit A, Otto R, et al. The Metabolic Requirements of Vinyasa Yoga. *Med Sci Sports Exerc.* 2003;35(5):S155.
128. Tran MD, Holly RG, Lashbrook J, Amsterdam EA. Effects of hatha yoga practice on the health-related aspects of physical fitness. *Prev Cardiol.* 2001;4(4):165-70.

129. Tracy BL, Hart CE. Bikram yoga training and physical fitness in healthy young adults. *J Strength Cond Res.* 2012.
130. Pullen PR, Nagamia SH, Mehta PK, et al. Effects of Yoga on Inflammation and Exercise Capacity in Patients With Chronic Heart Failure. *J Card Fail.* 2008;14(5):407-13.
131. Clay CC, Lloyd LK, Walker JLea. The metabolic cost of hatha yoga. *J Strength Cond Res.* 2005;19(3):604-10.
132. DiCarlo L, Sparling P, Hinson B, Snow T, Roszkopf L. Cardiovascular, metabolic, and perceptual responses to hatha yoga standing poses. *Med Exerc Nutr Health.* 1995;4:107-12.
133. Bowman AJ, Clayton RH, Murray A, et al. Effects of aerobic exercise training and yoga on the baroreflex in healthy elderly persons. *Eur J Clin Invest.* 1997;27(5):443-9.
134. Fenwick PBC, Donaldson S, Gillis L. Metabolic and EEG changes during transcendental meditation: an explanation. *Biol Psychiatry.* 1977;5(2):101-18.
135. Ramos-Jiménez A, Hernández-Torres RP, Wall-Medrano A, Muñoz-Daw MDJ, Torres-Durán PV, Juárez-Oropeza MA. Cardiovascular and metabolic effects of intensive Hatha Yoga training in middle-aged and older women from northern Mexico. *International journal of yoga.* 2009;2(2):49.
136. Throll DA. Transcendental meditation and progressive relaxation: Their physiological effects. *J Clin Psychol.* 1982;38(3):522-30.
137. Buranruk O, La Grow S, Ladawan S, et al. Thai yoga as an appropriate alternative physical activity for older adults. *J Complement Integr Med.* 2010;7(1).
138. Danucalov MĂD, SimĂmes RS, Kozasa EH, et al. Cardiorespiratory and metabolic changes during yoga sessions: The effects of respiratory exercises and meditation practices. *Appl Psychophysiol Biofeedback.* 2008;33(2):77-81.
139. Miyamura M, Nishimura K, Ishida K, Katayama K, Shimaoka M, Hiruta S. Is man able to breathe once a minute for an hour?: The effect of yoga respiration on blood gases. *Jpn J Physiol.* 2002;52(3):313-6.
140. Frostell C, Pande JN, Hedenstierna G. Effects of high-frequency breathing on pulmonary ventilation and gas exchange. *J Appl Physiol.* 1983;55(6):1854-61.
141. Vivekananda R. *Practical yoga psychology.* Munger, Bihar, India: Yoga Publications Trust; 2005.
142. Ramos-Jiménez A, Hernández-Torres R, Wall-Medrano A. Hatha Yoga Program Determinants on Cardiovascular Health in Physically Active Adult Women. *J Yoga Phys Therap.* 2011;1(103):2.
143. Joint F. *Energy and Protein Requirements: Report of a Joint FAO/WHO/UNU Expert Consultation;* World Health Organization; 1985.
144. Poehlman ET, Melby CL, Badylak SF. Resting metabolic rate and postprandial thermogenesis in highly trained and untrained males. *Am J Clin Nutr.* 1988 May 1, 1988;47(5):793-8.
145. Tremblay A, Fontaine E, Poehlman ET, et al. The effect of exercise-training on resting metabolic rate in lean and moderately obese individuals. *Int J Obes.* 1986;10(6):511-7.
146. Ray US, Mukhopadhyaya S, Purkayastha SS, et al. Effect of yogic exercises on physical and mental health of young fellowship course trainees. *Indian J Physiol Pharmacol.* 2001;45(1):37-53.
147. Ross A, Thomas S. The health benefits of yoga and exercise: a review of comparison studies. *J Altern Complement Med.* 2010;16(1):3-12. PubMed PMID: 2010546692. Language: English. Entry Date: 20100312. Revision Date: 20100312. Publication Type: journal article.
148. Van Dixhoorn J. Cardiorespiratory effects of breathing and relaxation instruction in myocardial infarction patients. *Biol Psychiatry.* 1998;49(1-2):123-35.
149. Clark ME, Hirschman R. Effects of paced respiration on anxiety reduction in a clinical population. *Biofeedback Self Regul.* 1990;15(3):273-84.
150. Kaushik RM, Kaushik R, Mahajan SK, Rajesh V. Effects of mental relaxation and slow breathing in essential hypertension. *Complement Ther Med.* 2006;14(2):120-6.
151. Joseph CN, Porta C, Casucci G, Casiraghi N, Maffei M, Rossi M, et al. Slow breathing improves arterial baroreflex sensitivity and decreases blood pressure in essential hypertension. *Hypertension.* 2005;46(4):714-8.
152. Ernst E. Breathing techniques - Adjunctive treatment modalities for asthma? A systematic review. *Eur Respir J.* 2000;15(5):969-72.
153. Cowen VS, Adams TB. Heart rate in yoga asana practice: A comparison of styles. *J Bodyw Mov Ther.* 2007;11(1):91-5.

154. Cowen VS, Adams TB. Physical and perceptual benefits of yoga asana practice: results of a pilot study. *J Bodyw Mov Ther.* 2005;9(3):211-9.
155. Wenger M, Bagchi B, .et.al. Studies on autonomic functions in practitioners of Yoga in India. *Behav Sci.* 1961;6:312 - 23.
156. WENGER MA, BAGCHI BK, ANAND BK. Experiments in India on "Voluntary" Control of the Heart and Pulse. *Circulation.* 1961 December 1, 1961;24(6):1319-25.
157. Kothari LK, Bordia A, Gupta OP. The yogic claim of voluntary control over the heart beat: An unusual demonstration. *Am Heart J.* 1973;86(2):282-4.
158. Benson H, Malhotra MS, Goldman RF, et al. Three case reports of the metabolic and electroencephalographic changes during advanced buddhist meditation techniques. *Behav Med.* 1990;16(2):90-5.
159. Lindholm P, Lundgren CE. The physiology and pathophysiology of human breath-hold diving. *J Appl Physiol.* 2009 January 2009;106(1):284-92.

